



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

system, 46; diseases of respiratory system, 43; diseases of digestive system, 83; diseases of urinary apparatus, 17; erysipelas, 3; malarial fevers, 26; plague, 17; septicæmia puerperal, 2; septicæmia other, 1; syphilis, 8; smallpox, 1; pulmonary tuberculosis, 61; tuberculosis, other, 2; enteric fever, 3; violent deaths not suicides, 5; stillborn, 27; other causes, 40; total, 543.

During the month there were reported 41 cases of plague, of which 6 were not confirmed. There were 12 deaths in residences and 5 in hospital, leaving 18 under treatment in hospital at close of month. All houses in which occurred cases of plague were disinfected, and 239 persons who had come in contact with cases were visited daily by sanitary inspectors.

There were 39 cases of smallpox, 24 male and 15 female, all removed to hospital, where there occurred 1 death.

During the month 334 houses were disinfected, 239 because of plague, 48 for tuberculosis, 43 for smallpox, 3 for enteric fever, and 1 for dysentery.

*Mortality during month of September, 1904—Plague and smallpox.*

Total number of deaths, 431. Causes: Alcoholism, 1; anæmia, 2; asthma, 1; beriberi, 29; Bright's disease, 3; cirrhosis of liver, 9; cancers, 7; cerebral hemorrhage and congestion, 18; convulsions, 15; diarrhea and dysentery, 42; diseases of circulatory system, 55; debility, congenital, 3; debility, senile, 16; erysipelas, 2; meningitis, 6; myelitis, 3; malarial fevers, 15; nephritis, 4; pneumonia, 9; puerperal hemorrhage, 1; plague, 10; plague suspect, 12; smallpox, 5; enteric fever, 1; septicæmia, puerperal, 1; septicæmia, other, 3; syphilis, 4; pulmonary tuberculosis, 47; tuberculosis, other, 3; tetanus, 6; still born, 31; other causes, 67.

There were 66 cases of smallpox, of which 5 died as stated above.

BRITISH HONDURAS.

*Report from Belize, fruit port.*

Acting Assistant Surgeon Carson reports as follows: Week ended October 13, 1904: Present officially estimated population, 8,500; number of deaths, 9; prevailing disease, malarial fever of mild type and phthisis pulmonalis; general sanitary condition of this port and the surrounding country during the week, fairly good.

Bills of health were issued to the following-named vessels:

Date.	Vessel.	Number of crew.	Number of passengers from this port.	Number of passengers in transit.	Pieces of baggage disinfected.
Oct. 7	Beverly .....	44	6	8	0
7	Falk .....	23	0	0	0
13	Colombia .....	18	0	0	0